



Technical Data Sheet

Recombinant Human MIP-1 alpha (CCL3) (rHu MIP-1 alpha (CCL3))

Human MIP-1 alpha /CCL3

Both MIP-1 alpha and MIP-1 beta are structurally and functionally related CC chemokines. They participate in the host response to invading bacterial, viral, parasite and fungal pathogens by regulating the trafficking and activation state of selected subgroups of inflammatory cells e.g. macrophages, lymphocytes and NK cells. While both MIP-1 alpha and MIP-1 beta exert similar effects on monocytes their effect on lymphocytes differ; with MIP-1 alpha selectively attracting CD8+ lymphocytes and MIP-1 beta selectively attracting CD4+ lymphocytes. Additionally, MIP-1 alpha and MIP-1 beta have also been shown to be potent chemoattractants for B cells, eosinophils and dendritic cells. Both human and murine MIP-1 alpha and MIP-1 beta are active on human and murine hematopoietic cells.

Catalog Number:	RC315-14
Source:	<i>Escherichia coli</i> .
Molecular Weight:	7.8 kDa protein containing 69 amino acid residues, including the four highly conserved cysteine residues present in CC chemokines.
Quantity:	5ug/20ug/1mg
Purity:	>96% by SDS-PAGE and HPLC analyses.
Biological Activity:	Determined by its ability to chemoattract human monocytes using a concentration range of 1.0-10.0 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2µm filtered concentrated (1.0mg/ml) solution in 20mM PB, pH 7.4, 100mM NaCl.
AA Sequence:	ASLAADTPTA CCFSYTSRQI PQNFIADYFE TSSQCSKPGV IFLTKRSRQV CADPSEEWVQ KYVSDLELSA
Endotoxin:	Less than 1EU/µg of rHu MIP-1 alpha /CCL3 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions.
Storage:	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long



Bio Basic Inc.

A world leader in serving science

term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. **Avoid repeated freeze/thaw cycles.**

Usage:

This material is offered by Bio Basic Inc. for research, laboratory or further evaluation purposes. **NOT FOR HUMAN USE.**