



Technical Data Sheet

Recombinant Human Keratinocyte Growth Factor-1 (rHu KGF-1)

Human Keratinocyte Growth Factor-1

Keratinocyte Growth Factor-1 (KGF-1/FGF-7) is one of 23 known members of the FGF family. All FGFs have two conserved cysteine residues and share 30 - 50% sequence identity at the amino acid level. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of variety of tissues, by promoting cellular proliferation and differentiation. KGF-1/FGF-7 is a mitogen factor specific for epithelial cells and keratinocytes and signals through FGFR 2b. KGF-1/FGF-7 plays a role in kidney and lung development, angiogenesis, and wound healing.

Catalog Number:	RC215-18
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 19.0 kDa, a single, non-glycosylated polypeptide chain containing 164 amino acids.
Quantity:	2ug/10ug/1000µg
Purity:	>96% by SDS-PAGE and HPLC analyses.
Biological Activity:	The biological activity was determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing KGF receptors yielding an ED ₅₀ <10ng/ml, corresponding to a specific activity of $\geq 1 \times 10^5$ units/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2µm filtered solution in 20mM PB, pH 8.0, 1M NaCl.
AA Sequence:	MCNDMTPEQMATNVNCSSPERHTRSVDYMEGGDIRVRRFCRTQWYLRIKRGKVKGT QEMKNNYNIMEIRTVAVGIVAIGVSEFYLAMNKEGKLYAKKECNEDCNFKELILENHNT YASAKWTHNGGEMFVALNOK GIPVRGKTK KEQKTAHFLP MAIT
Endotoxin:	Less than 1EU/µg of rHuKGF-1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq -20^\circ\text{C}$. Further dilutions should be made in appropriate buffered solutions.



Bio Basic Inc.

A world leader in serving science

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by Bio Basic Canada Inc for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.