



Technical Data Sheet

Recombinant Murine Interleukin-13 (rMu IL-13)

Mouse Interleukin-13

IL-13 is an immunoregulatory cytokine produced primarily by activated Th2 cells, and also by mast cells and NK cells. Targeted deletion of IL-13 in mice resulted in impaired Th2 cell development and indicated an important role for IL-13 in the expulsion of gastrointestinal parasites. IL-13 exerts anti-inflammatory effects on monocytes and macrophages and it inhibits the expression of inflammatory cytokines such as IL-1beta, TNF-alpha, IL-6 and IL-8. IL-13 has also been shown to enhance B cell proliferation and to induce isotype switching resulting in increased production of IgE. Blocking of IL-13 activity inhibits the pathophysiology of asthma. Human and murine IL-13 is cross-species reactive.

Catalog Number:	RC232-24
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 12.3 kDa, a single non-glycosylated polypeptide chain containing 111 amino acids.
Quantity:	2ug/10ug/1mg
Purity:	>97% by SDS-PAGE and HPLC analyses.
Biological Activity:	The ED50 as determined by the dose-dependent proliferation of TF-1 cells was < 4.0 ng/ml, corresponding to a specific activity of > 2.5 x 10 ⁵ units/mg.
Physical Appearance:	Sterile Filtered White Lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2µm filtered solution in PBS, pH 7.4.
AA Sequence:	MPVPRSVSLP LTLKELIEEL SNITQDQTPL CNGSMVWSVD LAAGGFCVAL DSLTNISNCN AIYRTQRILH GLCNRKAPTT VSSLPDTKIE VAHFITKLLS YTKQLFRHGP F
Endotoxin:	Less than 1EU/µg of rIL-13 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions.
Storage:	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.



Bio Basic Inc.

A world leader in serving science

Usage:

This material is offered by Bio Basic Inc. for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.