



Technical Data Sheet

Recombinant Human MCP-1/MCAF (rHu MCP-1/ MCAF (CCL2))

Human MCP-1/MCAF

CCL2, also known as monocyte chemoattract and activating factor (MCAF), was initially purified independently by two groups based on its ability to chemoattract monocytes. Subsequent to its cloning and sequencing, it became evident that this protein is also identical to the product of the human *JE* gene. The *JE* gene, originally identified in mouse fibroblasts, is a platelet-derived growth factor (PDGF)-inducible gene. The human CCL2 cDNA encodes a 99 amino acid residue precursor protein with a 23 residue hydrophobic signal peptide that is cleaved to generate the 76 residue mature protein. Natural CCL2 is heterogeneous in size due to the addition of O-linked carbohydrates and sialic acid residues. In addition to fibroblasts; tumor cells, smooth muscle cells, endothelial cells, and mononuclear phagocytes can also produce CCL2 either constitutively or upon stimulation by various stimuli. CCL2 is a member of the β (CC) subfamily of chemokines. Recently, the existence of MCP2 and MCP3 with 62% and 73% amino acid identity respectively, to CCL2 have been reported.

| | |
|-----------------------------|---|
| Catalog Number: | RC315-13 |
| Source: | <i>Escherichia coli</i> . |
| Molecular Weight: | 8.6 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids. |
| Quantity: | 5ug/20ug/1mg |
| Purity: | >96% by SDS-PAGE and HPLC analyses. |
| Biological Activity: | Measured by its ability to chemoattract human CCR2A transfected BaF3 mouse pro-B cells. The ED ₅₀ for this effect is typically 10-60 ng/mL. |
| Physical Appearance: | Sterile Filtered White lyophilized (freeze-dried) powder. |
| Formulation: | Lyophilized from a 0.2 μ m filtered concentrated (1.0mg/ml) solution in 20mM PB, pH 7.4, 100mM NaCl. |
| AA Sequence: | QPDAINAPVT CCYNFTNRKI SVQRLASYRR ITSSKCPKEA VIFKTIVAKE ICADPKQKWV QDSMDHLDKQ TQTPKT |
| Endotoxin: | Less than 1EU/ μ g of rHuMCP-1/ MCAF /CCL2 as determined by LAL method. |
| Reconstitution: | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at \leq -20°C. Further dilutions should be made in appropriate buffered solutions. |
| Storage: | This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable |



Bio Basic Inc.

A world leader in serving science

for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. **Avoid repeated freeze/thaw cycles.**

Usage:

This material is offered by Bio Basic Inc. for research, laboratory or further evaluation purposes. **NOT FOR HUMAN USE.**